arXiv:1007.2461v2 [math.MG] 16 Jul 2010

ON EDWARDS–CHILD'S INEQUALITY

YU-DONG WU*, ZHI-HUA ZHANG, AND ZHI-GANG WANG

Abstract. In this paper, by making use of one of Chen's theorems and the method of mathematical analysis, we refine Edwards–Child's inequality and solve a conjecture posed by Liu [6].

1. Introduction and Main Results

For $\triangle ABC$, let a, b, c denote the side-lengths, A, B, C the angles, p the semi-perimeter, R the circumradius and r the inradius, respectively.

Bottema *et al.* [1, pp.50, Theorem 5.8] once recorded the following so-called Edwards-Child's inequality in their monograph:

$$p \geqq 3\sqrt{3}\, r. \tag{1.1}$$

In 1999, Liu [5] considered the refinement of inequality (1.1) and posed the problem: Determine the best constant t for the inequality

$$a+b+c \geqq 6\sqrt{3}\, r + t \cdot \frac{r}{R} \cdot (R-2r) \Longleftrightarrow p \geqq 3\sqrt{3}\, r + \frac{t}{2} \cdot \left(1 - \frac{2r}{R}\right) r. \tag{1.2}$$

In the same year, Huang [4] solved the above problem and obtained the best constant

$$t = 6.829212418 \cdots .$$

In 2000, Liu [6] posed a conjecture which sharpened inequality (1.1) as follows:

Conjecture 1.1. (LBQ 67) *Prove or disprove the inequality:*

$$\sqrt{3}\, p \geqq 10r - r\left(\frac{2r}{R}\right)^5 \Longleftrightarrow p \geqq 3\sqrt{3}\, r + \frac{\sqrt{3}}{3}\left[1 - \left(\frac{2r}{R}\right)^5\right] r. \tag{1.3}$$

The main object of this paper is to prove and refine inequality (1.3), and we obtain the following results.

Theorem 1.1. *If* $\lambda \leqq f(x_1) \approx 5.977930729$*, then*

$$\sqrt{3}\, p \geqq 10r - r\left(\frac{2r}{R}\right)^{\lambda}, \tag{1.4}$$

the equality in (1.4) *is valid if and only if* $a : b : c = 1 : 1 : 1$ *with* $\lambda \leqq f(x_1)$ *or* $a : b : c = 2\left(x_1^2 - 3\right) : \left(x_1^2+3\right) : \left(x_1^2+3\right)$ *with* $\lambda = f(x_1)$*, where*

$$f(x) := \frac{\ln\left[-\frac{2(x^3-5x^2+15)}{x^2-3}\right]}{\ln\left[\frac{24(x^2-3)}{(x^2+3)^2}\right]} \qquad (3 < x < x_0), \tag{1.5}$$

Date: January 1, 2008.

2000 *Mathematics Subject Classification.* Primary 51M16; Secondary 26D15.

Key words and phrases. Edwards–Child's inequality; Best constant; Discriminant sequence; Sylvester's Resultant.

*Corresponding author, e-mail address: yudong.wu@hotmail.com.

$x_1 \approx 3.067873979 \in (3, x_0)$ *is the only positive real root of the equation* $f'(x) = 0$, *and* $x_0 \approx 4.113537611$ *is the maximal real root of the following equation:*

$$x^3 - 5x^2 + 15 = 0.$$

Theorem 1.2. *If* $k \leqq k_0$, *then for any triangle, we have*

$$p \geqq 3\sqrt{3}\ r + k\left[1 - \left(\frac{2r}{R}\right)^5\right] r, \tag{1.6}$$

the equality in (1.6) *is valid if and only if* $a : b : c = 1 : 1 : 1$ *with* $k \leqq k_0$ *or* $a : b : c = 2t_1 : 1 : 1$ *with* $k = k_0$, *where* $k_0 \approx 0.6898369707 \in \left(\frac{1}{2}, \frac{9}{13}\right)$ *is the root of the equation* $p_4(k) = 0$, $t_1 \approx 0.5194285605 \in \left(\frac{1}{2}, \frac{3}{5}\right)$ *is the root of the equation* $p_2(t) = 0$, *and*

$$\begin{aligned} p_4(k) = {} & 582076609134674072265625k^{40} \\ & - 347825698554515838623046875k^{38} \\ & + 100074581801891326904296875000k^{36} \\ & - 18656729921698570251464843750000k^{34} \\ & + 2406402165103435516357421875000000k^{32} \\ & - 223881811253170562744140625000000000k^{30} \\ & + 13463175828870323610839843750000000000k^{28} \\ & - 546554928845186341347265625000000000000k^{26} \\ & + 44602844570033253018161875000000000000000k^{24} \\ & - 5130242398470886015317438950000000000000000k^{22} \\ & + 805214383330095009369880969748800000000000000k^{20} \\ & + 8091804003905867976154901735852032000000000000k^{18} \\ & - 262076670696960781271382283372767764480000000000k^{16} \\ & + 483089080872113925346827615868453187944448000000000k^{14} \\ & + 76133654804831682593043073614469922630706135040000000k^{12} \\ & + 4656778296665388277933286028092539858012767959121920000k^{10} \\ & + 144475786697302680271016689740018636544036558347226316800k^{8} \\ & + 2305862812061518538327375046497265994061106944720616030208k^{6} \\ & + 14131109130840787698067340120866948829204788093469111353344k^{4} \\ & - 27802797644590317388762803455393575731200000000000000000000k^{2} \\ & + 9774552621457470122500000000000000000000000000000000000000, \end{aligned} \tag{1.7}$$

and

$$\begin{aligned} p_2(t) = {} & 91750400t^{20} - 857210880t^{19} + 3560046592t^{18} - 8616673280t^{17} \\ & + 13364330496t^{16} - 13704527872t^{15} + 9174425600t^{14} - 3719282688t^{13} \\ & + 689182720t^{12} + 53528576t^{11} - 36429312t^{10} + 533760t^{9} + 207104t^{8} \\ & + 75520t^{7} + 32352t^{6} + 11376t^{5} - 320t^{4} - 104t^{3} - 30t^{2} - 7t - 1. \end{aligned} \tag{1.8}$$

2. Preliminary Results

In order to prove our main results, we shall require the following three lemmas.

Lemma 2.1. (see [2, 3, 8]) *The homogeneous inequality* $p \geqq (>) f_1(R, r)$ *in triangle holds if and only if*

$$2(1+t)\sqrt{1-t^2} \geqq (>) f_1(1, 2t(1-t)), \tag{2.1}$$

where $t = \cos B = \cos C \in \left[\frac{1}{2}, 1\right)$. *And the homogeneous inequality* $p \leqq (<) f_1(R, r)$ *holds if and only if* (2.1) *is reversed, where* $t = \cos B = \cos C \in \left(0, \frac{1}{2}\right]$.

Lemma 2.2. (see [9, 10]) *Given a polynomial* $f(x)$ *with real coefficients*

$$f(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n,$$

if the number of the sign changes of the revised sign list of its discriminant sequence

$$\{D_1(f), D_2(f), \cdots, D_n(f)\}$$

is v*, then, the number of the pairs of distinct conjugate imaginary roots of* $f(x)$ *equals* v*. Furthermore, if the number of non-vanishing members of the revised sign list is* l*, then, the number of the distinct real roots of* $f(x)$ *equals* $l - 2v$.

Lemma 2.3. (see [10]) *Denote*

$$F(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n,$$

and

$$G(x) = b_0x^m + b_1x^{m-1} + \cdots + b_m.$$

If $a_0 \neq 0$ *or* $b_0 \neq 0$*, then the polynomials* $F(x)$ *and* $G(x)$ *have the common roots if and only if*

$$R(F,G) = \begin{vmatrix} a_0 & a_1 & a_2 & \cdots & a_n & 0 & \cdots & 0 \\ 0 & a_0 & a_1 & \cdots & a_{n-1} & a_n & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & a_0 & \cdots & \cdots & \cdots & a_n \\ b_0 & b_1 & b_2 & \cdots & \cdots & \cdots & \cdots & 0 \\ 0 & b_0 & b_1 & \cdots & \cdots & \cdots & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & b_0 & b_1 & \cdots & b_m \end{vmatrix} = 0,$$

where $R(F, G)$ *is Sylvester's Resultant of* $F(x)$ *and* $G(x)$.

3. The Proof of Theorem 1.1

Proof. By Lemma 2.1, we know that inequality (1.4) can be written as follows:

$$2\sqrt{3}(1+t)\sqrt{1-t^2} \geqq 20t(1-t) - 2t(1-t)[4t(1-t)]^{\lambda} \qquad \left(\frac{1}{2} \leqq t < 1\right). \tag{3.1}$$

If we let

$$x = \sqrt{\frac{3+3t}{1-t}},$$

then

$$t = \frac{x^2-3}{x^2+3} \qquad (x \geqq 3),$$

and inequality (3.1) is equivalent to

$$\left[\frac{24(x^2-3)}{(x^2+3)^2}\right]^{\lambda} \geqq -\frac{2(x^3-5x^2+15)}{x^2-3} \qquad (x \geqq 3). \tag{3.2}$$

We now split it into three cases to prove.

(i) When $x=3$, inequality (3.2) is an identity for $\lambda \in \mathbb{R}$ and $\triangle ABC$ is equilateral.

(ii) Clearly, if $-\frac{2(x^3-5x^2+15)}{x^2-3} \leqq 0$, or $x \geqq x_0 \approx 4.113537611$, then inequality (3.2) holds for $\lambda \in \mathbb{R}$.

(iii) When $3<x<x_0$, we have

$$0<\frac{24(x^2-3)}{(x^2+3)^2}<1,$$

and inequality (3.2) is equivalent to

$$\lambda \leqq f(x),$$

where $f(x)$ is given by (1.5). Calculating the derivative for $f(x)$, we get

$$f'(x)=\frac{x(x-3)(x+3)g(x)}{(x^2+3)(x^2-3)(x^3-5x^2+15)\ln^2\left[\frac{24(x^2-3)}{(x^2+3)^2}\right]}.$$

where

$$g(x):=x(x^2+3)\ln\left[\frac{24(x^2-3)}{(x^2+3)^2}\right]+2(x^3-5x^2+15)\ln\left[-\frac{2(x^3-5x^2+15)}{x^2-3}\right]$$
$$(3<x<x_0).$$

Let $f'(x)=0$, we obtain

$$g(x)=0. \tag{3.3}$$

Now, we prove equation (3.3) has only one real root on interval $(3,x_0)$.

For $g(x)$, we know that

$$g'(x)=3(x^2+1)\ln\left[\frac{24(x^2-3)}{(x^2+3)^2}\right]+2x(3x-10)\ln\left[-\frac{2(x^3-5x^2+15)}{x^2-3}\right],$$

$$\begin{aligned} g''(x)=&\frac{2x(x-3)(x+3)(5x^4+6x^3-60x^2-45)}{(x^3-5x^2+15)(x^2+3)(x^2-3)}+6x\ln\left[\frac{24(x^2-3)}{(x^2+3)^2}\right]\\ &+(12x-20)\ln\left[-\frac{2(x^3-5x^2+15)}{x^2-3}\right], \end{aligned}$$

$$\begin{aligned} g'''(x)=&\frac{2}{(x^3-5x^2+15)^2(x^2+3)^2(x^2-3)^2}\big(20x^{13}-113x^{12}-180x^{11}+1500x^{10}\\ &-540x^9-2349x^8+1620x^7-14256x^6-29160x^5+164025x^4+58320x^3\\ &-364500x^2-54675\big)+6\ln\left[\frac{24(x^2-3)}{(x^2+3)^2}\right]+12\ln\left[-\frac{2(x^3-5x^2+15)}{x^2-3}\right], \end{aligned}$$

and

$$g^{(4)}(x)=\frac{4xp(x)}{(x^3-5x^2+15)^3(x^2+3)^3(x^2-3)^3},$$

where

$$\begin{aligned} p(x)=&5x^{18}-78x^{17}+375x^{16}-306x^{15}-705x^{14}-4239x^{13}+13230x^{12}-16443x^{11}\\ &+213435x^{10}-1047330x^9+1356750x^8+4312764x^7-12141495x^6-2460375x^5\\ &+22766670x^4+4100625x^3-32805000x^2-14762250x+61509375. \end{aligned}$$

The revised sign list of the discriminant sequence of $p(x)$ is

$$[1,1,-1,1,1,-1,-1,-1,1,1,-1,-1,-1,1,1,-1,1,-1]. \tag{3.4}$$

So the number of the sign changes of the revised sign list of (3.4) is 9. By Lemma 2.2, we know that the number of the pairs of distinct conjugate imaginary roots of $p(x)$ are 9. Namely, $p(x)=0$ has no real root, then $p(x)>0$ for $x\in R$.

With $p(x)>0$ for $x\in R$, $g^{(4)}(x)<0$ for $x\in(3,x_0)$, then $g'''(x)$ is strictly decreasing on $(3,x_0)$. And for

$$g'''(3)=3>0 \quad \text{and} \quad g'''(4)=6\ln\left(\frac{312}{361}\right)-\frac{106269606}{61009}+12\ln\left(\frac{2}{13}\right)<0,$$

we conclude that $g'''(x)=0$ has only one real root $x_4\approx 3.016763142$ on $(3,x_0)$. So $g''(x)$ is strictly increasing on $(3,x_4)$, and decreasing on (x_4,x_0).

Similarly, we deduce that

$$g^{(j)}(x)=0 \qquad (0\leqq j\leqq 2)$$

have only one real root x_{j+1} $(0\leqq j\leqq 2)$ on $(3,x_0)$, respectively.

Therefore, we know that $f'(x)=0$ has only one real root $x_1\approx 3.067873979\in(3,x_0)$, and

$$\lambda_{\max}=f(x)_{\min}=f(x_1)\approx 5.977930729.$$

By Lemma 2.1, we find that the equality in (1.4) is valid when $\triangle ABC$ is an isoceles triangle, and if we let $b=c=1$, then

$$a=2\cos B=2\cos C=2t_1=\frac{2(x_1^2-3)}{x_1^2+3}.$$

Namely, the equality in (1.4) is valid if and only if

$$a:b:c=2\left(x_1^2-3\right):\left(x_1^2+3\right):\left(x_1^2+3\right)$$

with $\lambda=f(x_1)$. Furthermore, considering the proof above, we can easily obtain the required results of the equality in (1.4).

□

4. THE PROOF OF THEOREM 1.2

Proof. By Lemma 2.1, we know that inequality (1.6) is equivalent to

$$2(1+t)\sqrt{1-t^2}\geqq 6\sqrt{3}t(1-t)+k\cdot 2t(1-t)\left\{1-[4t(1-t)]^5\right\} \qquad \left(\frac{1}{2}\leqq t<1\right). \tag{4.1}$$

For $t=\frac{1}{2}$, (4.1) obviously holds and $\triangle ABC$ is equilateral. If $\frac{1}{2}<t<1$, then (4.1) is equivalent to

$$k\leqq h(t):=\frac{(1+t)\sqrt{1-t^2}-3\sqrt{3}t(1-t)}{t(1-t)\left\{1-[4t(1-t)]^5\right\}} \qquad \left(\frac{1}{2}<t<1\right). \tag{4.2}$$

Therefore, we get

$$\begin{aligned}h'(t)=&\big[5120t^{12}-14336t^{11}+1024t^{10}+30720t^9-25600t^8-10240t^7+19456t^6-6144t^5\\&+t+1+15360\sqrt{3}\sqrt{1-t^2}(1-t)^5t^6\big]/\big[t^2(1-t)(2t-1)^3(256t^8-1024t^7\\&+1472t^6-832t^5+80t^4+32t^3+12t^2+4t+1)^2\sqrt{1-t^2}\ \big].\end{aligned}$$

Let $h'(t)=0$, we obtain

$$\begin{aligned}&5120t^{12}-14336t^{11}+1024t^{10}+30720t^9-25600t^8-10240t^7+19456t^6\\&\quad-6144t^5+t+1+15360\sqrt{3}\sqrt{1-t^2}(1-t)^5t^6=0.\end{aligned} \tag{4.3}$$

It is easy to see that the roots of equation (4.3) must be the roots of the following equation:

$$p_2(t)(t+1)(2t-1)^3 = 0, \tag{4.4}$$

where $p_2(t)$ is given by (1.8).

The revised sign list of the discriminant sequence of $p_2(t)$ is

$$[1, 1, 1, -1, -1, 1, -1, -1, -1, 1, 1, 1, -1, 1, 1, 1, -1, -1, 1, 1]. \tag{4.5}$$

So the number of the sign changes of the revised sign list of (4.5) is 8, by Lemma 2.2, the equation $p_2(t) = 0$ has 4 distinct real roots. By using the function "fsolve()" in Maple 9.0, we can obtain the approximation of 2 distinct real roots $t_1 \approx 0.5194285605$ and $t_2 \approx 0.8281776966$ on open interval $\left(\frac{1}{2}, 1\right)$. But t_2 does not satisfy equation (4.3), which implies that it is an extraneous root. Hence, it follows that

$$h(t)_{\min} = h(t_1) = \frac{(1+t_1)\sqrt{1-t_1^2} - 3\sqrt{3}t_1(1-t_1)}{t_1(1-t_1)\{1-[4t_1(1-t_1)]^5\}} \approx 0.6898369707 \in \left(\frac{1}{2}, \frac{9}{13}\right). \tag{4.6}$$

We now prove $h(t_1)$ is the root of equation (3.1). We consider the nonlinear algebraic equation system as follows:

$$\begin{cases} p_2(t_1) = 0, \\ u^2 + t_1^2 - 1 = 0, \\ v^2 - 27t_1^2(1-t_1)^2 = 0, \\ (1+t_1)u - v - \left\{t_1(1-t_1)\left\{1-[4t_1(1-t_1)]^5\right\}\right\}k = 0, \end{cases} \tag{4.7}$$

or

$$\begin{cases} p_2(t_1) = 0, \\ p_3(t_1, k) = 0, \end{cases} \tag{4.8}$$

where $p_2(t)$ is given by (1.8) and

$$\begin{aligned} p_3(t,k) = {} & 68719476736k^4t^{42} - 1374389534720k^4t^{41} \\ & + 13022340841472k^4t^{40} - 77687368450048k^4t^{39} + 327083234426880k^4t^{38} \\ & - 1032364109070336k^4t^{37} + 2532355667394560k^4t^{36} - 4938039864328192k^4t^{35} \\ & + 7763189089435648k^4t^{34} - 9918303552143360k^4t^{33} + 10328492122046464k^4t^{32} \\ & - 8752260867686400k^4t^{31} + 5995600248045568k^4t^{30} - 3278892823478272k^4t^{29} \\ & + 1402218741760000k^4t^{28} - 453762372075520k^4t^{27} + 105664190873600k^4t^{26} \\ & - 16867120906240k^4t^{25} + 2458252738560k^4t^{24} - 944229580800k^4t^{23} \\ & + 65536k^2(-52 + 7344385k^2)t^{22} - 1441792k^2(-24 + 119075k^2)t^{21} \\ & + 4587520k^2(8441k^2 - 34)t^{20} - 65536k^2(65905k^2 - 6238)t^{19} \\ & + 20480k^2(-33404 + 4115k^2)t^{18} - 32768k^2(2431k^2 - 22973)t^{17} \\ & + 2048k^2(29895k^2 - 262244)t^{16} - 2048k^2(12617k^2 - 113240)t^{15} \\ & + 256k^2(-196132 + 23549k^2)t^{14} - 2560k^2(39k^2 - 1232)t^{13} \\ & - 512k^2(1929 + 71k^2)t^{12} - 256k^2(-1576 + 55k^2)t^{11} - 16k^2(109k^2 - 2660)t^{10} \\ & - 64k^2(51k^2 - 1274)t^9 + 40k^2(-296 + 7k^2)t^8 + 8k^2(-788 + 11k^2)t^7 \\ & + k^2(25k^2 - 3076)t^6 + 2k^2(-400 + 3k^2)t^5 + k^2(k^2 - 166)t^4 \\ & - 20k^2t^3 + (49 - 2k^2)t^2 + 14t + 1. \end{aligned}$$

It's easy to see that $h(t_1)$ is also the solution of the nonlinear algebraic equation system (4.8). By Lemma 2.3, we get

$$R(p_2, p_3) = mp_4(k)p_5(k) = 0, \tag{4.9}$$

where

$$\begin{aligned} m =\,&5769822192075864843731878595793471609238717138319092113989119658\\ &3608446361959417370721935453787479709740599154674963543698529626\\ &6124545032425816659873003224168956415237423809191615340664942700\\ &6115796999896656147107615727318549627125299675136000000000, \end{aligned}$$

$p_4(k)$ is given by (1.7), and

$$\begin{aligned} p_5(k) = {}& 886065557319729123264551162719726562 5k^{40}\\ &- 16912080685346637403103522956371307373046875k^{38}\\ &+ 543574184102733722721203416585922241210937 5000k^{36}\\ &- 1022146637691011812806702821850776672363281250000k^{34}\\ &+ 110744571814857445274894881859313964843750000000000k^{32}\\ &- 93693894608112829735297577655916583251953125000000000k^{30}\\ &+ 2824410316091286848110397826839951184570312500000000000k^{28}\\ &- 1038042584540888321508236786385886072604187500000000000000k^{26}\\ &+ 73340535725547628521708894311600819307265000000000000000000k^{24}\\ &- 265177139233592420219133919392417723053323929920000000000000000k^{22}\\ &+ 11791633182359761811369142031608045030413197651840000000000000000k^{20}\\ &+ 190075444754710412043307126977386168137413366964111360000000000k^{18}\\ &- 1092700953333652491900906640137222767517014418741343404032000000000k^{16}\\ &- 77504079403992396391275778207617305822909294158004250227507200000000k^{14}\\ &- 1280167121239702258044546676265726571686977622628855665332846592000000k^{12}\\ &+ 28596041041659567331601090256258283539207054207485483280129169817600000k^{10}\\ &+ 110267783320883416462258578945595318603200831554078609670205562316390400k^{8}\\ &+ 745324132711516724121423152749040579524897407250894331064162322304991232k^{6}\\ &- 7375168578676180534548979250039227170865907645609839475147982859847860224k^{4}\\ &- 802076741103259008414533593438831455813452680864661504000000000000000000000k^{2}\\ &+ 395739228446715359134735466442056102500000000000000000000000000000000000000. \end{aligned} \tag{4.10}$$

The revised sign list of the discriminant sequence of $p_5(k)$ is

$$\begin{gathered} [1, 1, 1, 1, -1, 1, 1, -1, 1, 1, 1, 1, -1, 1, -1, 1, 1, 1, 1, -1, 1, 1, 1, -1, 1, 1, 1, 1,\\ 1, -1, -1, -1, 1, 1, -1, 1, 1, 1, 1, 1]. \end{gathered} \tag{4.11}$$

So the number of the sign changes of the revised sign list of (4.11) is 16, by Lemma 2.2, the equation $p_5(k) = 0$ has 8 distinct real roots. And making use of the function "realroot()"[7] in Maple 9.0, we know that the equation $p_5(k) = 0$ has 8 distinct real roots in the following intervals:

$$\left[2, \frac{5}{2}\right], \quad \left[\frac{5}{2}, 3\right], \quad [4, 8], \quad [1024, 2048], \quad \left[-\frac{5}{2}, -2\right], \quad \left[-3, -\frac{5}{2}\right], \quad [-8, -4], \quad [-2048, -1024].$$

Therefore, the equation $p_5(k) = 0$ has no real root on interval $\left(\frac{1}{2}, \frac{9}{13}\right)$.

From (4.6), we find that $h(t_1)$ is the root of the equation $p_4(k) = 0$. Namely, $h(t_1)$ is the root of the equation (1.7).

Furthermore, by the similar arguments of the last part of the proof of Theorem 1.1, we can easily get the necessary and sufficient conditions on the equality in (1.6).

We thus complete the proof of Theorem 1.2. □

Acknowledgements

The present investigation was supported, in part, by the *Hunan Provincial Natural Science Foundation* under Grant 05JJ30013 and the *Scientific Research Fund of Hunan Provincial Education Department* under Grant 05C266 of People's Republic of China.

References

[1] O. Bottema, R. Ž. Djordević, R. R. Janić, D. S. Mitrinović and P. M. Vasić, *Geometric Inequalities*, Wolters-Noordhoff Publishing, Groningen, The Netherlands, 1969.

[2] S.-L. Chen, *A new method to prove one kind of inequalities—equate substitution method*, Fujian High-School Mathematics, **3** (1993), 20–23. (in Chinese)

[3] S.-L. Chen, *The simplified method to prove inequalities in triangle*, Studies of Inequalities, Tibet People's Press (2000), 3–8. (in Chinese)

[4] B.-C. Huang, *Solutions of 13 problems*, Commun. Studies Inequal. **6** (1999), 17–22. (in Chinese)

[5] B.-Q. Liu, *182 geometric inequalities problems in triangle*, Commun. Studies Inequal. **6** (1999), 48–63. (in Chinese)

[6] B.-Q. Liu, *110 interesting inequalities problems*, Studies of Inequalities, Tibet People's Press (2000), 398. (in Chinese)

[7] D.-M. Wang. *Selected lectures in symbolic computation.* Tsinghua University Press, Beijing, **8** (2003), 110–114. (in Chinese)

[8] Y.-D. Wu, *The best constant for a geometric inequality*, J. Inequal. Pure Appl. Math. **6** (2005), Art. 111. (electronic)

[9] L. Yang, X.-R. Hou and Z.-B. Zeng, *A complete discrimination system for polynomials*, Sci. China (Ser. E), **39** (1996), 628–646.

[10] L. Yang, J.-Z. Zhang and X.-R. Hou, *Nonlinear algebraic equation system and automated theorem proving*, Shanghai Scientific and Technological Education Press (1996), 23–25. (in Chinese)

(Y.-D. Wu) Department of Mathematics, Zhejiang Xinchang High School, Shaoxing, Zhejiang 312500, P. R. China.
E-mail address: `yudong.wu@hotmail.com`

(Z.-H. Zhang) Department of Mathematics, Shili Senior High School in Zixing, Chenzhou, Hunan 423400, P. R. China.
E-mail address: `zxzh1234@163.com`

(Z.-G. Wang) School of Mathematics and Computing Science, Changsha University of Science and Technology, Changsha, Hunan 410076, P. R. China.
E-mail address: `zhigwang@163.com`